
Stamp of the practice:

Standard Operating Procedure (SOP)







	1	T	
WHAT	WHEN	HOW	BY MEANS OF
Cleaning and disinfection of the tray	Prior to the new preparation of the cleaning solution	Wear protective gloves, an apron impermeable to water + protective glasses! > empty the tray > rinse the tray with water > if necessary clean with a suitable brush > wipe out the tray completely with a disinfection wipe	Plastic brush MinutenWipes, PlastiSept- Wipes eco MaxiWipes soaked with MinutenSpray-classic or PlastiSept eco
Preparation of cleaning solution	 1 x per working day at visible contamination (e. g. clouding, coagulation, streaking) 	Wear protective gloves, an apron impermeable to water + protective glasses! > fill in tap water > add concentrate > mix well	Suitable tray Cleaning solution 1. warm tap water 2. TarClean 5 – 20 % depending on degree of soiling

Page 1 of 3 As at: 07.01.2016

WHAT	WHEN	HOW	BY MEANS OF
Pre-cleaning of dental workpieces (e. g. prostheses, bridges, dental splints, surgical masks etc.)	immediately after removal from the mouth	Wear protective gloves, an apron impermeable to water + protective glasses! During transport to the sterilization area to be protected against contamination rinse impression, dental workpieces under running water	Suitable water
Cleaning of dental workpieces (e. g. prostheses, bridges, dental splints, surgical masks etc.)	 prior to the transport to the laboratory after receipt from the laboratory 	Wear protective gloves, an apron impermeable to water + protective glasses! bubble-free deposit impressions and dental workpieces to be covered completely with the solution contact time x min.	TarClean depending on degree of soiling 5 – 20 % in the tray max. 30 minutes, in the ultrasonic device approx. 5 minutes
Disinfection see work instruction "Manual cleaning and disinfection of instruments in immersion bath (wet/chemical)"	 after the cleaning prior to the transport to the laboratory after receipt from the laboratory 	Wear protective gloves, an apron impermeable to water + protective glasses! bubble-free deposit impressions and dental workpieces to be covered completely with the solution contact time x min.	BIB-forte <i>eco</i> 0,5 % (60 min.) 3 % (30 min.) 4 % (5 min. ultrasonic d.) (tap water)

Page 2 of 3 As at: 07.01.2016

WHAT	WHEN	HOW	BY MEANS OF
Final step	After the disinfection	Wear protective gloves, an apron impermeable to water + protective glasses! > rinse under running water > blow dry	suitable water compressed air WL-Blow
Transport		Wear protective gloves, an apron impermeable to water + protective glasses! > packed to the external laboratory direct processing in the own laboratory	

Instruction by:	taken note of:	
Practice operator	Staff	

Page 3 of 3 As at: 07.01.2016